

Atmospheric pressure as a natural climate regulator for a terrestrial planet with a biosphere

King-Fai Li¹, Kaveh Pahlevan, Joseph L. Kirschvink, and Yuk L. Yung

Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125

Edited by Norman H. Sleep, Stanford University, Stanford, CA, and approved April 10, 2009 (received for review September 24, 2008)

Lovelock and Whitfield suggested in 1982 that, as the luminosity of the Sun increases over its life cycle, biologically enhanced silicate weathering is able to reduce the concentration of atmospheric carbon dioxide (CO₂) so that the Earth's surface temperature is maintained within an inhabitable range. As this process continues, however, between 100 and 900 million years (Ma) from now the CO₂ concentration will reach levels too low for C₃ and C₄ photosynthesis, signaling the end of the solar-powered biosphere. Here, we show that atmospheric pressure is another factor that adjusts the global temperature by broadening infrared absorption lines of greenhouse gases. A simple model including the reduction of atmospheric pressure suggests that the life span of the biosphere can be extended at least 2.3 Ga into the future, more than doubling previous estimates. This has important implications for seeking extraterrestrial life in the Universe. Space observations in the infrared region could test the hypothesis that atmospheric pressure regulates the surface temperature on extrasolar planets.

evolution | Gaia | greenhouse | habitability

The Earth is inhabitable largely because its atmosphere contains 2 principal greenhouse gases (water vapor and CO₂) that serve as a blanket to trap heat emitted from the surface; these regulate the global mean temperature within the range in which life can evolve. If such an atmosphere were absent, the global mean temperature would be only -18°C (1), and life, which relies on liquid water, would not exist. The standard model of stellar evolution (2) suggests that the solar luminosity was only 70% of the current value when the Sun formed, and has been steadily increasing to its current value over the past 4.6 billion years. This implies that the global mean temperature should have been increasing with time. However, the geological record indicates that our planet has stayed well within the inhabitable range of ≈ 0 – 50°C since the Archean (3), aside from a few possible “Snowball Earth” events (4, 5). Climate models also suggest that the partial pressure of CO₂ at the surface (P_{CO_2}) 3,500 Ma ago was as large as 7,000 Pa and has declined to the preindustrial value of 28 Pa (6). Had P_{CO_2} remained unchanged at the level of 7,000 Pa, the present-day surface temperature would have been much more than 50°C , the upper limit for most eukaryotic life. Walker et al. (7) suggested abiotic carbonate–silicate weathering as a mechanism that reduces the atmospheric CO₂, but it was Lovelock and Whitfield (3) who first proposed biologically enhanced silicate weathering as the primary mechanism regulating the temperature throughout the history of the Earth. The model proposed by Lovelock and Whitfield (3) suggested that P_{CO_2} would drop to ≈ 15 Pa, the lower limit for photosynthesis by C₃ plants, in ≈ 100 Ma, implying the end of the photosynthetic biosphere. This idea was further developed by Caldeira and Kasting (8) with detailed calculations including C₄ plants that can use CO₂ at pressures as low as 1 Pa. As the solar luminosity increases, the partial pressure of CO₂ should drop to 15 Pa in 0.5 billion years or gigaannum (Ga) and to 1 Pa in 0.9 Ga, eventually reaching a runaway greenhouse (9) that renders the Earth's surface uninhabitable. In their model, a saturated atmosphere was assumed, which allows the positive feedback by water vapor as the Earth's surface warms up. More elaborate

models that include biological and geodynamic processes have also been developed (10, 11), but estimates for the life span of the biosphere remain at ≈ 1 Ga.

All of these previous studies focused on the greenhouse effect due to direct absorption by atmospheric species such as water vapor and CO₂, whose radiative forcings are $\approx 80\text{ Wm}^{-2}$ and $\approx 30\text{ Wm}^{-2}$, respectively. However, atmospheric pressure also plays a critical role in the greenhouse effect through broadening of the infrared absorption lines of these gases by collisional interaction with other molecules (mainly N₂ and O₂ in the present atmosphere) (1). In other words, if the total atmospheric pressure were lower, the climate forcing of greenhouse gases would be smaller, the magnitude of the greenhouse effect would be less, and the global mean temperature would drop (see Fig. 2 and later discussions). If, via natural or artificial processes, it were possible to reduce the Earth's atmospheric pressure, the life span of the biosphere would be extended.

Geochemical and geobiological processes may have varied atmospheric pressure throughout the history of the Earth (12). Biology regulates overall atmospheric pressure in 2 related but distinct ways. First, the $\approx 2 \times 10^4$ Pa of oxygen in the present atmosphere is maintained directly by oxygenic photosynthesis: photosystem-II, operating in green plants, algae, and cyanobacteria. With the possible exception of O₂ produced by the sudden release of peroxides accumulated in the ice caps of a Snowball Earth (5), no other mechanism—biotic or abiotic—is known to be capable of maintaining even trace amounts of this gas in the Earth's atmosphere (see ref. 4 for a recent discussion).

The second pressure component in the Earth's atmosphere that interacts with biology is the reservoir of gaseous N₂; at nearly 78% volume mixing ratio (vmr), it has the largest influence on pressure broadening of the greenhouse gas absorption bands. The atmospheric nitrogen budget is ultimately tied to the oceanic primary productivity and the subsequent burial of organic material in the sediments. In the present ocean, organic matter is incorporated into living organisms in the stoichiometric Redfield ratio of 106:16:1 for C, N, and P (13). Thus, the C/N ratio is ≈ 6.6 . But when the organic carbon is buried in the marine sediments, its C/N ratio is much larger because N, an essential nutrient, has been stripped for recycling (see, e.g., chapter 11 of ref. 14). There are measurements of nitrogen in sediments, hard crustal rocks, and the mantle (15–17): The estimated amounts in the atmosphere, crust, and mantle are 2.8 , 0.4 ± 0.2 , and 5.5 ± 1.9 , respectively, in units of 10^{20} mol of N (15). These measurements show that it is possible to sequester on the order of an atmosphere of N₂ in the crust and mantle. The higher N/rock ratios seen in the data are believed to be associated with biology.

Thus, natural processes may have caused variations in atmospheric N₂ in the history of the Earth, a result that is

Author contributions: K.-F.L., K.P., J.L.K., and Y.L.Y. designed research; K.-F.L. performed research; K.-F.L. contributed new reagents/analytic tools; K.-F.L. analyzed data; and K.-F.L., K.P., J.L.K., and Y.L.Y. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed at: 1200 East California Boulevard, MC 150-21, Pasadena, CA 91125. E-mail: kfl@gps.caltech.edu.




Fig. 1. U.S. Standard Tropical Atmosphere profiles used in this work. Note that 1 mb = 100 Pa.

testable by measurements. For instance, basalt vesicularity has been used to infer atmospheric paleopressure and hence paleoelevation by using a standard atmosphere that is the same as the present one (18). Recently, measurement of atmospheric N_2 in the Archean period has also been proposed (29).

Model and Results

Given the potential for biological control of both oxygen and nitrogen, we cannot assume that the abundance of either of these gases has remained constant over geologic time. Here, we present a model to estimate how much the life span of the biosphere might be extended by regulating atmospheric pressure. We emphasize that the model to be presented is maximally simplified and that it aims only to estimate the first-order effect due to the pressure broadening. We shall discuss the shortcomings of the model in the final paragraph of this section. We assume that during the next 900 million years, the atmospheric

pressure is kept constant at the present value (10^5 Pa). Following Caldeira and Kasting (8), as the solar luminosity increases, the surface temperature rises to 300 K, and P_{CO_2} drops logarithmically to 1 Pa. At this time, the column abundances of CO_2 and other trace gases including water vapor remain constant such that C_4 plants can subsist. In our model, the atmospheric pressure is lowered to maintain the surface temperature T_{surf} at 300 K. For a given solar luminosity, the effective temperature T_{eff} of the Earth at radiative equilibrium can be obtained from,

$$\sigma T_{eff}^4 = \frac{(1-a)}{4} S, \quad [1]$$

where $a \approx 0.3$ is the planetary albedo and S is the solar luminosity measured at 1 AU, which is a function of time given by ref. 8

$$S(t) = S_0 (1 - 0.38t/4.55 \text{ Ga})^{-1}, \quad [2]$$

where $S_0 = 1,368 \text{ Wm}^{-2}$ is the current value of the solar luminosity (19). This incoming solar energy must be balanced by the outgoing longwave radiation, defined through the Stefan-Boltzmann law (1, 20):

$$\sigma T_{eff}^4 = \int_0^\infty \pi I_\nu(T, P, \rho_{GH}) d\nu, \quad [3]$$

where $\sigma = 5.67 \times 10^{-8} \text{ WmK}^{-4}$ is the Stefan-Boltzmann constant. I_ν is the outgoing radiation measured at the top of the atmosphere at wave number ν , which is a function of the vertical profiles of temperature $T(z)$, pressure $P(z)$, and the mixing ratios $\rho_{GH}(z)$ of the greenhouse gases; it is related to atmospheric pressure due to the pressure broadening effect described above. Thus, given the solar luminosity $S(t)$ at a particular time t and the profiles $\rho_{GH}(z)$ and $T(z)$ with $T_{surf} = T(z=0)$, one aims to solve for the function $P(z)$ such that Eqs. 1 and 3 are equal. For simplicity, we assume that $P(z) = \beta P_{ref}(z)$, i.e., the pressure profile is scaled as a whole, for some reference pressure profiles such that the problem is reduced to solving for the scalar constant β .




Fig. 2. Absorption spectra at the top of the atmosphere in the present-day Earth for $\beta = 1$ (blue) and 0.2 atm (orange), showing the effect of pressure broadening. The spectral resolution is 20 cm^{-1} . The major species responsible for absorption are labeled. (Inset) The total radiative forcing due to the absorption by greenhouse gases as a function of β .




Fig. 3. The evolution of surface pressure and temperature and the partial pressure of CO₂ in the model as the solar luminosity evolves in the next 2.8 Ga from present. Before 0.9 Ga (dashed line), the quantities evolve as described in Caldeira and Kasting (8). After 0.9 Ga, the surface temperature and the partial pressure of CO₂ at the surface (P_{CO_2}) are kept constant while the atmospheric pressure is lowered to compensate the temperature increase due to the increase in solar luminosity (see *Models and Results*). At ≈ 2.3 Ga (dash-dotted line), runaway greenhouse occurs, signaling the ultimate fate of the biosphere.

I_v is calculated from the MODTRAN radiative transfer code (21) by using the atmospheric profiles from the U.S. Standard Tropical Atmosphere data (22), including $P_{\text{ref}}(z)$ (the reference profile), $\rho_{\text{GH}}(z)$ and $T(z)$, where the surface temperature $T_{\text{surf}} \approx 300$ K (Fig. 1). We assume that $T(z)$ remains constant after 0.9 Ga. Fig. 2 illustrates the effect of pressure broadening on the absorption features due to the major greenhouse gases. The absorption spectra have been averaged to a resolution of 20 cm^{-1} . It is evident that the amount of absorption is reduced at lower pressure. The area under the absorption spectra can be integrated to give the total radiative forcing, which is shown in the Fig. 2 *Inset* as a function of β . As the solar luminosity evolves, we solve for the scaling factor β such that the equality between Eqs. 1 and 3 holds and that the energy balance is achieved:

$$\frac{(1-a)}{4}S(t) = \int_0^\infty \pi I_v [T(z), \beta(t)P_{\text{ref}}(z), \beta(t)^{-1}\rho_{\text{GH}}(z)] dv \quad [4]$$

Here, $\beta(t)$, which is time dependent as is $S(t)$, is the only parameter to be determined. Except for ozone, which is photochemically produced from the ambient oxygen gas whose concentration scales with the atmospheric pressure, $\rho_{\text{GH}}(z)$ is inversely scaled by $\beta(t)^{-1}$ such that the total columns of major greenhouse gases such as H₂O and CO₂ are kept constant.

The derived evolution of the surface pressure, given by $\beta(t)P_{\text{ref}}(z=0)$, is shown in Fig. 3. It drops to 3,500 Pa at ≈ 2.5 Ga, when the atmosphere consists mostly of water vapor. When the atmosphere becomes aquatic, most plants and other aerobic forms of life will be wiped out, marking the end of the aerobic biosphere. Only bacteria that are capable of carrying out anaerobic metabolisms can survive and continue the life processes. Thus, 2.5 Ga serves as a biological lower limit for the life span




Fig. 4. The emission spectra of the Earth's atmosphere corresponding to the 3 stages shown in Fig. 3. Blue: Stage A with 330 ppm CO₂ under 1 atmospheric pressure; red: Stage B with 10 ppm CO₂ under 1 atmospheric pressure; and, green: Stage C with 10 ppm CO₂ under 0.08 atmospheric pressure.

of the biosphere. This is a coarse estimate in that we have neglected the collision-induced absorption of water vapor. It is natural to ask whether the Earth's surface pressure will in fact decrease according to the previous condition. As pointed out by Walker (p 118 in ref. 23), nitrogen fixation by biology acts to naturally draw down the nitrogen content of the atmosphere through the process of organic matter burial. We can estimate an upper limit on the efficacy of this process as follows. The carbon isotope record (24) reveals that organic matter has accounted for $\approx 20\%$ of the carbon sinks at the Earth's surface in the last 3.5 Ga. Considering that a vast reservoir (≈ 60 bars CO₂ equivalent) of carbonates exists in sediments, this implies a reservoir of organic matter (≈ 10 bars CO₂ equivalent) including nitrogen, which, with a Redfield ratio of C/N ≈ 6 , would imply 1 bar of N₂ having been removed from the atmosphere over the past 4 Ga, a rate that is comparable with that needed to offset the effects of increasing solar luminosity.

However, it is difficult to extend the life span of the biosphere much beyond 2 Ga physically because the runaway greenhouse limits the stability of the atmosphere against loss of water vapor from the planet. In the present atmosphere, loss of water is mitigated by the cold tropopause. Therefore, even though the mixing ratio of H₂O is $\approx 1\%$ near the surface, this value drops to $\approx 10^{-6}$ near the tropopause, effectively trapping the H₂O in the troposphere. However, this cold trap would fail if the atmosphere were composed of $\approx 40\%$ H₂O. In this case the moist lapse rate would be insufficient to cause cold trapping at the tropopause; H₂O would be abundant in the upper atmosphere, where it would be dissociated by UV radiation, followed by escape to space (9). This is probably the explanation of why Venus is so dry compared with Earth, having lost an ocean of H₂O via failure of the cold trap. Referring to Fig. 3, $T_{\text{surf}} \approx 27^\circ\text{C}$ implies $P_{\text{H}_2\text{O}} \approx 3,500$ Pa. The runaway greenhouse threshold P_{RG} may be estimated from $P_{\text{H}_2\text{O}}/P_{\text{RG}} \approx 0.4$, from which we obtain $P_{\text{RG}} \approx 8,900$ Pa or 0.08 atm for the atmosphere to be stable against escape. The pressure must not fall below P_{RG} , which according to our model occurs 2.3 Ga from present. (dash-dotted line in Fig. 3).

In the above calculations, we have neglected biological or geological feedback mechanisms like those in refs. 10 and 11). For example, as the Earth's mantle cools down in time, the volcanic source of atmospheric CO₂ may accordingly drop (10). In the absence of feedbacks, this would imply a lower level of atmospheric CO₂ in steady state than the constant value of 10

Pa assumed in Fig. 3 after 0.9 Ga. This difficulty arises because we have considered the fall of CO₂ and the pressure loss separately in different time domains, whereas in reality, they will operate simultaneously, the modeling of which requires additional research.

Discussion

Such a prolonged life span for the aerobic biosphere has implications for the search for life on extrasolar planets and the question of intelligent life in the Universe (25). The presence of oxygen-derived ozone (O₃) in the atmospheres of planets is considered to be a biomarker that enables remote identification of the presence of life. Because the Earth has already expressed this signature for 2.3 Ga, the continued persistence of the photosynthetic biosphere for 2.3 Ga into the future implies that the Earth will be identifiable as an inhabited planet for nearly half the total lifetime of the Sun, an important point to consider in the search for life on extrasolar planets.

In general, inert gases besides N₂ may provide significant pressure broadening in the atmospheres of exoplanets. Origin scenarios allow for a wide range of atmospheric masses and compositions for forming planets. Thus, given the amount of greenhouse gases in the atmosphere, the type of star the planet is next to and the orbital distance of the planet from the star, one can anticipate that the parameter that determines whether a planet is habitable is atmospheric pressure. Indeed, this is the major reason why present-day Mars is not inhabitable.

Remote sensing measurements using low-resolution infrared spectra have provided convincing evidence of the habitability of extrasolar planets (26) and the search continues. In the Earth's atmosphere, the dominant absorption band in the infrared region at $\approx 700\text{ cm}^{-1}$ is due to stratospheric CO₂, whose bandwidth is $\approx 200\text{ cm}^{-1}$ ($\Delta\nu/\nu \approx 0.28$). Thus, this CO₂ band is the most conspicuous signature to examine the pressure broadening effect in the atmosphere of an Earth-like extrasolar planet. Fig.

4 shows the IR spectra of the Earth's atmosphere corresponding to the 3 stages shown in Fig. 3: (A) The present atmosphere (330 ppm CO₂ at 1 atm); (B) The atmosphere 0.9 Ga after present as predicted by Caldeira and Kasting (8) (10 ppm CO₂ at 1 atm); and, (C) The atmosphere ≈ 2.3 Ga after present as predicted by the present authors (10 ppm CO₂ at 0.08 atm). In Stage A, the atmosphere is simulated with a water vapor abundance half that in the other 2 to account for a low vapor pressure due to the 10 K difference in the surface temperatures. With a resolution of $\Delta\nu \approx 100\text{ cm}^{-1}$ ($\Delta\nu/\nu \approx 0.14$), the absorption band at $\approx 700\text{ cm}^{-1}$ is clearly resolved. The absorption is reduced by half when CO₂ drops from 330 to 10 ppm; it is reduced by another half when the pressure drops by a factor of 10. Differences in the water vapor absorption are also apparent. However, these absorption features cannot be distinguished from those in the current atmosphere (e.g., the Spitzer/IRS instrument has a resolution $\Delta\nu/\nu \approx 0.5$ near 700 cm^{-1}) unless the spectral resolution is at least twice as fine as the bandwidth of CO₂. This poses a challenge for future space technology.

Model Summary

The MODTRAN radiative transfer code has been used in this study (21). It has a moderate spectral resolution for wavelengths extending from thermal infrared to UV regions and it uses spherical geometry to compute the radiances. It solves the radiative transfer equations by the discrete ordinate method (27). We simulate the outgoing longwave radiation at the top of the atmosphere. The 1 cm^{-1} statistical band model (1) is used, where the molecular band parameters are derived from the HITRAN database. The Voigt equivalent width is directly evaluated from the exact expression for finite spectral bands (28).

ACKNOWLEDGMENTS. We thank D. Feldman, V. Natraj, R.-L. Shia, and C. Parkinson for reading the manuscript. We also thank Prof. Norm Sleep and 2 anonymous reviewers for their insightful suggestions. K.F.L. and Y.L.Y. were supported by National Aeronautics and Space Administration Grant NNG06GF33G and the Virtual Planetary Laboratory at the California Institute of Technology.

- Goody RM, Yung YL (1989) *Atmospheric Radiation: Theoretical Basis* (Oxford Univ Press, Oxford) 2nd Ed.
- Gough DO (1981) Solar interior structure and luminosity variations. *Sol Phys* 74(1):21–34.
- Lovelock JE, Whitfield M (1982) Life-span of the biosphere. *Nature* 296(5857):561–563.
- Kopp RE, Kirschvink JL, Hilburn IA, Nash CZ (2005) Was the Paleoproterozoic Snowball Earth a biologically-triggered climate disaster? *Proc Natl Acad Sci USA* 102:11131–11136.
- Liang MC, Hartman H, Kopp RE, Kirschvink JL, Yung YL (2006) Production of hydrogen peroxide in the atmosphere of a Snowball Earth and the origin of oxygenic photosynthesis. *Proc Natl Acad Sci USA* 103(50):18896–18899.
- Owen T, Cess RD, Ramanathan V (1979) Enhanced CO₂ greenhouse to compensate for reduced solar luminosity on early Earth. *Nature* 277(5698):640–642.
- Walker JCG, Hays PB, Kasting JF (1981) A negative feedback mechanism for the long-term stabilization of Earth's surface temperature. *J Geophys Res Oceans* 86(C10):9776–9782.
- Caldeira K, Kasting JF (1992) The life-span of the biosphere revisited. *Nature* 360(6406):721–723.
- Ingersoll A (1969) Runaway greenhouse—A history of water on Venus. *J Atmos Sci* 26(6):1191–1198.
- Franck S, et al. (2000) Reduction of biosphere life span as a consequence of geodynamics. *Tellus B* 52(1):94–107.
- Lenton TM, von Bloh W (2001) Biotic feedback extends the life span of the biosphere. *Geophys Res Lett* 28(9):1715–1718.
- Hart MH (1978) Evolution of atmosphere of Earth. *Icarus* 33(1):23–39.
- Emerson SR, Hedges JI (2008) *Chemical Oceanography and the Marine Carbon Cycle* (Cambridge Univ Press, Cambridge, UK).
- Burdige DJ (2006) *Geochemistry of Marine Sediments* (Princeton Univ Press, Princeton).
- Marty B, Dauphas N (2003) The nitrogen record of crust-mantle interaction and mantle convection from Archean to present. *Earth Planet Sci Lett* 206(3–4):397–410.
- Papineau D, Mojzsis SJ, Karhu JA, Marty B (2005) Nitrogen isotopic composition of ammoniated phyllosilicates: Case studies from Precambrian metamorphosed sedimentary rocks. *Chem Geol* 216(1–2):37–58.
- Watenphul A, Wunder B, Heinrich W (2009) High-pressure ammonium-bearing silicates: Implications for nitrogen and hydrogen storage in the Earth's mantle. *Am Miner* 94(2–3):283–292.
- Sahagian D, Prussevitch A (2007) Paleoelevation measurement on the basis of vesicular basalts. *Mineral Geochem* 66:195–213.
- Woods T, et al. (2006) Solar Irradiance Variability during the SORCE Mission. *The Earth Observer* 18(4):8–14.
- Ackerman SA, Frey RA, Smith WL (1992) Radiation budget studies using collocated observations from advanced very high-resolution radiometer, high-resolution infrared sounder/2, and earth radiation budget experiment instruments. *J Geophys Res Atmos* 97(D11):11513–11525.
- Kneizy FX, et al. (1996) *The MODTRAN 2/3 Report and LOWTRAN 7 Model* (Phillips Lab, Geophysics Directorate, PL/GPOS, Hanscom AFB, Waltham, MA).
- Anderson GP, Clough SA, Kneizy FX, Chetwynd JH, Shettle EP (1986) AFGL atmospheric constituent profiles (0–120 km) in *Tech Rep AFGL-TR-86-0110*, AFGL (OPI) (Phillips Lab, Hanscom AFB, Waltham, MA).
- Walker JCG (1977) *Evolution of the Atmosphere* (Macmillan, New York), p 318.
- Schidlowski M (1987) Application of stable carbon isotopes to early biochemical evolution on Earth. *Annu Rev Earth Planet Sci* 15:47–72.
- Beichman CA, Woolf NJ, Lindensmith CA, eds (1999) *The Terrestrial Planet Finder* (JPL Publication, Pasadena, CA).
- Tinetti G, et al. (2007) Water vapour in the atmosphere of a transiting extrasolar planet. *Nature* 448(7150):169–171.
- Stamnes K, Tsay SC, Wiscombe W, Jayaweera K (1988) Numerically stable algorithm for discrete-ordinate-method radiative-transfer in multiple-scattering and emitting layered media. *Appl Opt* 27(12):2502–2509.
- Berk A, Bernstein LS, Robertson DC (1989) *MODTRAN: A Moderate Resolution Model for LOWTRAN 7*, Report GL-TR-89-0122 (Phillips Lab, Geophysics Directorate, Hanscom AFB, Waltham, MA).
- Schirber M (2008) Measuring the weight of ancient air. *Astrobiology Magazine*. Available at <http://www.astrobio.net/news/>. Accessed October 06, 2008.